

Seasonal Timing of Diseases – Fruit Crops



Fruit Diseases	Fruit crop species affected	Seedlings	Roots	Branches/Canes	Foliage	Flowers	Fruit
Disease Name		APRIL	MAY	JUNE	JULY	AUGUST	SEPTEMBER
Entomosporium Leaf and Berry Spot	Saskatoon berry	Solid line from April to May	Solid line from May to June	Solid line from June to August	Solid line from June to August	Solid line from July to August	Solid line from August to September
Mummyberry / Black Rot	Saskatoon berry		Solid line from May to June	Solid line from June to July			
Saskatoon berry / Juniper Rust	Saskatoon berry (juniper = secondary host)		Solid line from May to June	Solid line from June to July	Solid line from July to August	Solid line from August to September	
Blackleaf / Witches' Broom	Saskatoon berry	Solid line from April to May	Solid line from May to June	Solid line from June to August	Solid line from July to August	Solid line from August to September	Solid line from September to October
Fireblight	All plants within Rosaceae (saskatoon berry, apple, etc.), caneberries		Solid line from May to June	Solid line from June to July	Solid line from July to August	Solid line from August to September	Solid line from September to October
Botrytis Grey Mold	Strawberry, raspberry, etc.	Solid line from April to May	Solid line from May to June	Solid line from June to August	Solid line from July to August	Solid line from August to September	Solid line from September to October
Powdery Mildew	Strawberry, Saskatoon berry, raspberry, black currants		Solid line from May to June	Solid line from June to July	Solid line from July to August	Solid line from August to September	Solid line from September to October
Pythium Root Rot ¹	Strawberry, raspberry, etc.	Solid line from April to May	Solid line from May to June	Solid line from June to August	Solid line from July to August	Solid line from August to September	Solid line from September to October
SOLID LINE = DIRECT	DASHED LINE = INDIRECT	1 – whenever younger plants are present		2 – Depends on weather		3 – Depends on crop stage	



Seasonal Timing of Diseases – Fruit Crops



Fruit Diseases	Fruit crop species affected	Seedlings	Roots	Branches/Canes	Foliage	Flowers	Fruit
Disease Name		APRIL	MAY	JUNE	JULY	AUGUST	SEPTEMBER
Cane Blight	Caneberries (raspberries, etc.)			●————●	●————●	●————●	●————●
Spur Blight	Caneberries (raspberries, etc.)			●————●	●————●	●————●	●————●
Common Leaf Spot	Strawberry		●————●		●————●	●————●	●————●
Black Knot	Prunus spp. (cherries, chokecherries, etc.)	●————●		●————●	●————●		●————●
Brown Rot / Cherry Blossom Blight	Prunus spp. (cherries, chokecherries, etc.)	●————●	●————●	●————●	●————●	●————●	●————●
Cytospora Canker	Saskatoon berry; other crops (different spp.)	●————●		●————●	●————●		●————●
Root Rots ²	Raspberry, etc.	●————●	●————●	●————●	●————●	●————●	●————●
Black Root Rot ²	Strawberry	●————●	●————●	●————●	●————●	●————●	●————●
Verticillium Wilt ^{2,3}	Strawberry, raspberry, other fruit	●————●	●————●	●————●	●————●	●————●	●————●
SOLID LINE = DIRECT	DASHED LINE = INDIRECT	1 – whenever younger plants are present		2 – Depends on weather		3 – Depends on crop stage	

